

Gevrey vectors of multi-quasi-elliptic systems

Chikh BOUZAR and Rachid CHAILI

ABSTRACT. We show that the multi-quasi-ellipticity is a necessary and sufficient condition for the property of elliptic iterates to be hold for multi-quasi-homogenous differential operators.

1. Introduction

The aim of this work is to prove the property of elliptic iterates for multi-quasi-elliptic systems of differential operators in generalized Gevrey spaces $G^{\mathcal{F},s}(\Omega)$, where $\mathcal{F}$ denotes the Newton's polyhedron of the system $(P_j)_{j=1}^N$. The property of elliptic iterates for the system $(P_j)_{j=1}^N$ in the generalized Gevrey classes $G^{\mathcal{F},s}(\Omega)$ means the inclusion

$$G^s\left(\Omega, (P_j)_{j=1}^N\right) \subset G^{\mathcal{F},s}(\Omega)$$

Let $P_j(x, D) = \sum_{\alpha} a_{j\alpha}(x_0) D^{\alpha}$, $j = 1, \dots, N$, denoted $(P_j)_{j=1}^N$, be linear differential operators with C^{∞} coefficients in a open subset Ω of $\mathbb{R}^n$.

DEFINITION 1. Fix $x_0 \in \Omega$, we define the Newton's polyhedron of the system $(P_j)_{j=1}^N$ at the point x_0 , noted $\mathcal{F}(x_0)$, as the convex hull of the set $\{0\} \cup \{\alpha \in \mathbb{N}^n, \exists j \in \{1, \dots, N\}; a_{j\alpha}(x_0) \neq 0\}$. A Newton's polyhedron $\mathcal{F}$ is said regular if there exists a finite set $Q(\mathcal{F}) \subset (\mathbb{R}_+^*)^n$ such that

$$\mathcal{F} = \bigcap_{q \in Q(\mathcal{F})} \{q \in \mathbb{R}_+^n, \langle \alpha, q \rangle \leq 1\}$$

If $S(\mathcal{F})$ designs the set of vertices of $\mathcal{F}$, we put

$$V(\xi) = \sum_{\alpha \in S(\mathcal{F})} |\xi^{\alpha}|,$$

1991 *Mathematics Subject Classification.* Primary 35B65, 35H10; Secondary 35N10.

Key words and phrases. Differential operators. Multi-quasi-ellipticity. Gevrey vectors. Gevrey spaces.

and

$$\begin{aligned} k(\alpha, \mathcal{F}) &= \inf \{t > 0, t^{-1}\alpha \in \mathcal{F}\}, \quad \alpha \in \mathbb{R}_+^n \\ \mu_j(\mathcal{F}) &= \max_{q \in Q(\mathcal{F})} q_j^{-1}, \quad j = 1, \dots, n \\ \mu(\mathcal{F}) &= \max_{1 \leq j \leq n} \mu_j(\mathcal{F}) \\ \theta(\mathcal{F}) &= \left(\frac{\mu(\mathcal{F})}{\mu_1(\mathcal{F})}, \dots, \frac{\mu(\mathcal{F})}{\mu_n(\mathcal{F})} \right) \end{aligned}$$

DEFINITION 2. Let $\mathcal{F}$ be a regular polyhedron and $s \in \mathbb{R}_+$, we denote $G^{\mathcal{F},s}(\Omega)$ the space of $u \in C^\infty(\Omega)$ such that $\forall H$ compact of Ω , $\exists C > 0, \forall \alpha \in \mathbb{N}^n$,

$$(1.1) \quad \sup_H |D^\alpha u| \leq C^{|\alpha|+1} [\Gamma(\mu(\mathcal{F})k(\alpha, \mathcal{F}) + 1)]^s$$

REMARK 1. We can take $\sup_H |D^\alpha u|$ or $\|D^\alpha u\|_{L^2(H)}$ in the definition, according to Sobolev imbedding theorems.

DEFINITION 3. The system $(P_j)_{j=1}^N$ is said multi-quasi-elliptic in Ω if

- 1) $\mathcal{F}(x) = \mathcal{F}$ does not depend of $x \in \Omega$
- 2) $\mathcal{F}$ is regular
- 3) $\forall x \in \Omega, \exists C > 0, \exists R \geq 0, \forall \xi \in \mathbb{R}^n, |\xi| \geq R,$

$$V(\xi) \leq C \sum_{j=1}^N |P_j(x, \xi)|$$

DEFINITION 4. Let $(P_j)_{j=1}^N$ be a system of linear differential operators satisfying the conditions 1) and 2) of definition 3 and $s \in \mathbb{R}_+$, we denote $G^s(\Omega, (P_j)_{j=1}^N)$ the space of $u \in C^\infty(\Omega)$ such that $\forall H$ compact of Ω , $\exists C > 0, \forall l \in \mathbb{N}, 1 \leq i_l \leq N$,

$$(1.2) \quad \|P_{i_1} \dots P_{i_l} u\|_{L^2(H)} \leq C^{l+1} (l!)^{s\mu(\mathcal{F})}$$

The aim of this work is to show the following theorem.

THEOREM 1. Let Ω be an open subset of $\mathbb{R}^n$, $\sigma > s \geq 1$ and $(P_j(x, D))_{j=1}^N$ be a system of linear differential operators with $G^{\theta(\mathcal{F}),\sigma}(\Omega)$ coefficients, then

$$(P_j)_{j=1}^N \text{ is multi-quasi-elliptic in } \Omega \iff G^s(\Omega, (P_j)_{j=1}^N) \subset G^{\mathcal{F},s}(\Omega)$$

For differential operators with constant coefficients we have shown in [4] a more general result.

2. Sufficient condition

Instead of $Q(\mathcal{F}), k(\mathcal{F}, \alpha), \mu(\mathcal{F}), \mu_j(\mathcal{F}), \theta(\mathcal{F})$ we write, respectively, $Q, k(\alpha), \mu, \mu_j, \theta$. Denote $\mathcal{K} = \{k = k(\alpha) : \alpha \in \mathbb{N}^n\}$ and ω any open subset of $\mathbb{R}^n$. If $u \in C^\infty(\omega)$ and $k \in \mathcal{K}$, define

$$|u|_{k,\omega} = \sum_{k(\alpha)=k} \|D^\alpha u\|_{L^2(\omega)},$$

if $u \in C_0^\infty(\mathbb{R}^n)$ we write $|u|_k$. For $\rho > 0$, we denote $B_\rho = \left\{ x \in \mathbb{R}^n, \sum_{j=1}^n x_j^{2\mu_j/\mu} < \rho^2 \right\}$.

Let $(P_j)_{j=1}^N$ be a system of linear differential operators with coefficients defined in an open neighborhood Ω of the origin satisfying the following conditions

- i) The system $(P_j)_{j=1}^N$ is multi-quasi-elliptic in Ω .
- ii) The coefficients $a_{j\alpha} \in G^{\theta,s}(\Omega)$, $\forall \alpha \in \mathbb{N}^n, \forall j \in \{1, \dots, N\}$.

We define for $j = 1, \dots, N$, and $h \in \mathbb{N}$,

$$P_j^h(x, D) = \underbrace{P_j(x, D) \circ \dots \circ P_j(x, D)}_{h \text{ times}}.$$

From the multi-quasi-ellipticity of the system $(P_j)_{j=1}^N$ and following the proof of the lemma 3.4 of [7], we easily obtain

LEMMA 1. *There exists $\rho_0 > 0$, $\forall \varepsilon \in]0, \frac{1}{v(n)}[$, ($v(n)$ denote the number of elements of $\mathcal{K} \cap [0, n]$), we have*

$\exists C_1 > 0, \exists C_2(\varepsilon) > 0, \forall \delta \in]0, 1[, \forall \rho > 0, B_{\rho+\delta} \subset B_{\rho_0}, \forall u \in C^\infty(B_{\rho_0}), \forall p \geq n$,

$$(2.1) \quad \begin{aligned} |u|_{p+1, B_\rho} &\leq C_1 \left(\sum_{j=1}^N |P_j^n(x, D)u|_{p-n+1, B_{\rho+\delta}} + \varepsilon |u|_{p+1, B_{\rho+\delta}} + (\varepsilon\delta)^{-n\mu} |u|_{p-n+1, B_{\rho+\delta}} \right. \\ &\quad \left. + \sum_{h=0}^p \left(\frac{(p+1)!}{h!} \right)^{s\mu} C_2(\varepsilon)^{p+1-h} |u|_{h, B_{\rho+\delta}} \right), \end{aligned}$$

and for $p \leq n$, we have

(2.2)

$$|u|_{p+1, B_\rho} \leq C_1 \left(\sum_{j=1}^N |P_j^n(x, D)u|_{p-n+1, B_{\rho+\delta}} + \varepsilon |u|_{p+1, B_{\rho+\delta}} + (\varepsilon\delta)^{-(p+1)\mu} |u|_{0, B_{\rho+\delta}} \right)$$

Let $\lambda > 0$ and $R > 0$, for $p \in \mathbb{N}$, we set

$$\sigma_p(u, \lambda) = \sigma_p(u, \lambda, R) = (p!)^{-s\mu} \lambda^{-p} \sup_{R/2 \leq \rho < R} (R - \rho)^{p\mu} |u|_{p, B_\rho}$$

LEMMA 2. *Let ρ_0 be as in the lemma1 and let $0 < R < 1$ such that $\overline{B_R} \subset B_{\rho_0}$, then $\exists \lambda_0 > 0$ (λ_0 depends only of R and $(P_j)_{j=1}^N$), $\forall u \in C^\infty(B_{\rho_0}), \forall \lambda \geq \lambda_0, \forall p \geq n$,*

(2.3)

$$\sigma_{p+1}(u, \lambda) \leq [(p-n+2) \dots (p+1)]^{-s\mu} \sum_{j=1}^N \sigma_{p-n+1}(P_j^n u, \lambda) + \sum_{h=0}^p \sigma_h(u, \lambda),$$

and for $p \leq n-1$,

$$(2.4) \quad \sigma_{p+1}(u, \lambda) \leq (p+1)!^{-s\mu} \sum_{j=1}^N \sigma_0(P_j^{p+1} u, \lambda) + \sigma_0(u, \lambda)$$

PROOF. Let $p \geq n$, multiplying both sides of inequality (2.1) by $(p+1)!^{-s\mu} \lambda^{-p-1} (R - \rho)^{p\mu}$, taking $\delta = \frac{R-\rho}{p-n+2}$ and then passing to the sup over $\rho \in [R/2, R]$, we obtain

$$\sigma_{p+1}(u, \lambda) \leq C_1 (I_1 + \varepsilon I_2 + \varepsilon^{-n\mu} I_3 + I_4),$$

where

$$\begin{aligned}
I_1 &= \sum_{j=1}^N \frac{(R-\rho)^{(p+1)\mu}}{(p+1)!^{s\mu} \lambda^{p+1}} |P_j^n u|_{p-n+1, B_{\rho+\delta}} \\
&\leq \sum_{j=1}^N \left(\frac{(p-n+1)!}{(p+1)!} \right)^{s\mu} \frac{e^\mu}{\lambda^n} \sigma_{p-n+1}(P_j^n u, \lambda) \\
I_2 &= \frac{(R-\rho)^{(p+1)\mu}}{(p+1)!^{s\mu} \lambda^{p+1}} |u|_{p+1, B_{\rho+\delta}} \leq (2^n e)^\mu \sigma_{p+1}(u, \lambda) \\
I_3 &= \frac{(R-\rho)^{(p+1)\mu}}{(p+1)!^{s\mu} \lambda^{p+1}} \delta^{-n\mu} |u|_{p-n+1, B_{\rho+\delta}} \leq \frac{e^\mu}{\lambda^n} \sigma_{p-n+1}(u, \lambda) \\
I_4 &= \frac{(R-\rho)^{(p+1)\mu}}{(p+1)!^{s\mu} \lambda^{p+1}} \sum_{h=0}^p \left(\frac{(p+1)!}{h!} \right)^{s\mu} C_2(\varepsilon)^{p+1-h} |u|_{h, B_{\rho+\delta}} \\
&\leq \frac{e^\mu C_2(\varepsilon)}{\lambda} \sum_{h=0}^p \left(\frac{C_2(\varepsilon)}{\lambda} \right)^{p-h} \sigma_h(u, \lambda)
\end{aligned}$$

By a suitable choice of ε , we find

$$\begin{aligned}
\sigma_{p+1}(u, \lambda) &\leq \left(\frac{(p-n+1)!}{(p+1)!} \right)^{s\mu} \frac{\tilde{C}_1}{\lambda^n} \sum_{j=1}^N \sigma_{p-n+1}(P_j^n u, \lambda) + \frac{\tilde{C}_2}{\lambda^n} \sigma_{p-n+1}(u, \lambda) \\
&\quad + \frac{\tilde{C}_3}{\lambda} \sum_{h=0}^p \left(\frac{\tilde{C}_4}{\lambda} \right)^{p-h} \sigma_h(u, \lambda)
\end{aligned}$$

It suffices to take $\lambda_0 = \tilde{C}_1 + \tilde{C}_2 + \tilde{C}_3 + \tilde{C}_4$ to get (2.3). For the inequality (2.4) we multiply both sides of inequality (2.2) by $\frac{(R-\rho)^{(p+1)\mu}}{(p+1)!^{s\mu} \lambda^{p+1}}$, take $\delta = \frac{R-\rho}{2}$ and then we follow the same procedure as for (2.3) ■

LEMMA 3. Let ρ_0, R and λ_0 be as in lemma 2, then $\forall u \in C^\infty(B_{\rho_0})$, $\forall \lambda \geq \lambda_0$, $\forall p \in \mathbb{N}$,

$$(2.5) \quad \sigma_{p+1}(u, \lambda) \leq 2^{p+1} \sigma_0(u, \lambda) + \sum_{l=1}^{p+1} 2^{p+1-l} C_{p+1}^l \frac{1}{(l!)^{s\mu}} \sum_{\substack{1 \leq j \leq l \\ 1 \leq i_j \leq N}} \sigma_0(P_{i_1} \dots P_{i_l} u, \lambda)$$

PROOF. It's obtained by recurrence over p as in lemma 3.2 of [3]. ■

Our first result is the following theorem.

THEOREM 2. Let Ω be an open subset of $\mathbb{R}^n$, $s \geq 1$ and let $(P_j(x, D))_{j=1}^N$ be a system of linear differential operators with $G^{\theta, s}(\Omega)$ coefficients, then

$$(P_j)_{j=1}^N \text{ is multi-quasi-elliptic in } \Omega \Rightarrow G^s(\Omega, (P_j)_{j=1}^N) \subset G^{\mathcal{F}, s}(\Omega)$$

PROOF. It is sufficient to check (1.1) in a neighborhood of every point x of Ω . Let x be the origin, then there exist ρ_0, λ_0 and R such that the lemmas hold.

Let $u \in G^s(\Omega, (P_j)_{j=1}^N)$, then there exists $C_1 > 0$ such that

$$\sigma_0(P_{i_1} \dots P_{i_l} u, \lambda_0) \leq C_1^{l+1} (l!)^{s\mu}, \quad \forall l \in \mathbb{N},$$

hence from (2.5), we obtain

$$\sigma_{p+1}(u, \lambda_0) \leq C_1 (2 + NC_1)^{p+1}, \quad \forall p \in \mathbb{N},$$

which gives

$$(2.6) \quad |u|_{p+1, B_{R/2}} \leq (p+1)!^{s\mu} C_2^{(p+1)\mu+1}, \quad \forall p \in \mathbb{N}$$

Let $k \in \mathcal{K}$, if $k \geq n-1$, then $k = p+r$, where $p = [k-n] + 1$ and $r \in [n-1, n[$. The interpolation inequality from [7, lemme 2.3] gives

$$|u|_{k, B_{R/2}} \leq \varepsilon \left(|u|_{p+n, B_{R/2}} + |u|_{p+n-1, B_{R/2}} + \sum_{\substack{t \in \mathcal{K} \\ p \leq t < p+n-1}} |u|_{t, B_{R/2}} \right) + C\varepsilon^{-\frac{r}{n-r}} |u|_{p, B_{R/2}}$$

If we set $\varepsilon = \left(\frac{\Gamma(k+1)}{\Gamma(p+n+1)} \right)^{s\mu}$, then we obtain, with (2.6),

$$|u|_{k, B_{R/2}} \leq C_2^{k\mu+1} (\Gamma(k+1))^{s\mu} (2 + CC_3) + \sum_{\substack{t \in \mathcal{K} \\ p \leq t < p+n-1}} |u|_{t, B_{R/2}},$$

where C_3 is the constant of the inequality

$$\left(\frac{\Gamma(p+n+1)}{\Gamma(p+r+1)} \right)^{\frac{r}{n-r}} \leq C_3^{1/\mu} \frac{\Gamma(p+r+1)}{\Gamma(p+1)}, \quad p \in \mathbb{N}, \quad r \in \mathcal{K} \cap [0, n[$$

Applying again the above interpolation inequality to $\sum_{\substack{t \in \mathcal{K} \\ h \leq t < h+n-1}} |u|_{t, B_{R/2}}$, $h =$

$0, \dots, p$, we find

$$|u|_{k, B_{R/2}} \leq C_2^{k\mu+1} (\Gamma(k+1))^{s\mu} (2 + CC_3) \sum_{j=0}^p v(n)^j,$$

where $v(n)$ is the number of elements of $\mathcal{K} \cap [0, n[$.

Similarly, if $k \leq n-1$, we find

$$|u|_{k, B_{R/2}} \leq C_2^{k\mu+1} (\Gamma(k+1))^{s\mu} (1 + CC_3)$$

Hence for all $k \in \mathcal{K}$, we have

$$(2.7) \quad |u|_{k, B_{R/2}} \leq \tilde{C}^{k\mu+1} (\Gamma(k+1))^{s\mu}$$

By the imbedding theorem of anisotropic Sobolev spaces, if $l \in \mathcal{K}$ and $l > \frac{2}{k(e)}$,

where $k(e) = \max_{q \in Q} \sum_{j=1}^n q_j$, then $\exists C > 0, \forall \alpha \in \mathbb{N}^n$,

$$\sup_{B_{R/2}} |D^\alpha u| \leq C |D^\alpha u|_{l, B_{R/2}},$$

consequently from (2.7) we get the estimate (1.1) ■

As a consequence of this theorem, we obtain a result of Gevrey hypo-ellipticity for multi-quasi-elliptic systems.

COROLLARY 1. *Under the assumptions of theorem 2, the following propositions are equivalent*

- i) $u \in \mathcal{D}'(\Omega)$, $P_j u \in G^{\mathcal{F}, s}(\Omega)$, $\forall j = 1, \dots, N$.
- ii) $u \in G^{\mathcal{F}, s}(\Omega)$.

3. Necessary condition

In this section we prove the reciprocal of the theorem 2. For this aim we need a characterization of the multi-quasi-ellipticity for the system $(P_j(x, D))_{j=1}^N$, known in the case of a scalar operator in [5].

PROPOSITION 1. *A system $(P_j)_{j=1}^N$, satisfying 1) and 2) of Definition 2, is multi-quasi-elliptic in Ω if and only if for any $x \in \Omega$, $\forall q \in Q$,*

$$\sum_{j=1}^N |P_{jq}(x, \xi)| \neq 0, \quad \forall \xi \in \mathbb{R}^n, \xi_1 \dots \xi_n \neq 0,$$

where P_{jq} is the q -quasi-homogenous part of P_j , i.e.

$$P_{jq}(x, \xi) = \sum_{\langle \alpha, q \rangle = 1} a_{j\alpha}(x) \xi^\alpha$$

THEOREM 3. *Let Ω be an open subset of $\mathbb{R}^n$ and $P_j(x, D)$, $j = 1, \dots, N$, be differential operators with $G^{\theta, \sigma}(\Omega)$ coefficients, if $s > \sigma \geq 1$, then*

$$G^s(\Omega, (P_j)_{j=1}^N) \subset G^{\mathcal{F}, s}(\Omega) \Rightarrow (P_j)_{j=1}^N \text{ is multi-quasi-elliptic in } \Omega$$

PROOF. Assume that the system $(P_j)_{j=1}^N$ is not multi-quasi-elliptic, then there exists $x_0 \in \Omega$, $\exists q \in Q$ and $\exists \xi_0 \in S^{n-1}$, $\xi_{0,1} \dots \xi_{0,n} \neq 0$, such that

$$(3.1) \quad P_{jq}(x_0, \xi_0) = 0, \quad \forall j = 1, \dots, N$$

We shall construct a function $u \in G^s(\Omega, (P_j)_{j=1}^N)$ and $u \notin G^{\mathcal{F}, s}(\Omega)$, which contradicts the hypothesis. Choose ε such that

$$0 < \varepsilon \leq \frac{\mu(s - \sigma)}{2\mu s - \sigma} < \frac{1}{2} \text{ and } \varepsilon \leq \min_{\langle \beta, q \rangle < 1} \mu(1 - \langle \beta, q \rangle),$$

and put $\eta = \frac{1 - \varepsilon/\mu}{\mu s}$.

let $\delta > 0$ such that the ball $B_0 = B(x_0, 2\delta)$ be relatively compact in Ω , and let $\varphi \in G^{q, \sigma\mu}(\mathbb{R}^n)$ with compact support in $B(0, 2\delta)$ and $\varphi(x) \equiv 1$ in $B(0, \delta)$. The desired function is

$$u(x) = \int_1^{+\infty} \varphi[r^{\varepsilon q}(x - x_0)] e^{-r^\eta} e^{i \langle x - x_0, r^q \xi_0 \rangle} dr,$$

where $r^q x = (r^{q_1} x_1, r^{q_2} x_2, \dots, r^{q_n} x_n)$.

As in [6], we easily show, for sufficiently large m , that

$$|D^{m\alpha} u(x_0)| > \frac{1}{2\eta} |\xi_0^{m\alpha}| \Gamma\left(\frac{\langle m\alpha, q \rangle + 1}{\eta}\right)$$

If we choose α such that $k(\alpha) = \langle \alpha, q \rangle$, we obtain

$$|D^{m\alpha} u(x_0)| > \frac{c(\mu)^{\langle m\alpha, q \rangle}}{2\eta} |\xi_0^{m\alpha}| [\Gamma(\mu k(m\alpha) + 1)]^{\frac{1}{\mu\eta}},$$

and, since $\frac{1}{\mu\eta} > s$, then $u \notin G^{\mathcal{F}, s}(U_0)$ for any neighborhood U_0 of x_0 .

Let show that $u \in G^s \left(\Omega, (P_j)_{j=1}^N \right)$. Since the coefficients of P_j are in $G^{\theta, \sigma}(\Omega) \subset G^{q, \sigma \mu}(\Omega)$, thus $\exists M > 0, \forall \alpha \in \mathbb{Z}_+^n, \forall \beta \in \mathbb{Z}_+^n, \forall x \in B_0, \forall r \geq 1, \forall j = 1, \dots, N$,

$$(3.2) \quad \left| \left(D_x^\beta P_j^{(\alpha)} \right) (x, r^q \xi_0) \right| \leq M^{|\beta|+1} [\Gamma(< \beta, q > + 1)]^{\sigma \mu} r^{1- < \alpha, q >}$$

In the other hand in view of (3.1) we have

$$\forall \delta > 0, \exists C_1 > 0, \forall r \geq 1, \forall x \in \Omega, |x - x_0| < 2\delta r^{-\varepsilon/\mu}, \forall j = 1, \dots, N,$$

$$(3.3) \quad |P_j(x, r^q \xi_0)| \leq C_1 r^{1-\varepsilon/\mu}$$

Indeed

$$\begin{aligned} |P_j(x, r^q \xi_0)| &\leq |P_{jq}(x, r^q \xi_0)| + \sum_{< \beta, q > < 1} r^{< \beta, q >} |a_{j\beta}(x)| \left| \xi_0^\beta \right| \\ &\leq r |P_{jq}(x, \xi_0) - P_{jq}(x_0, \xi_0)| + \sum_{< \beta, q > < 1} r^{< \beta, q >} |a_{j\beta}(x)| \\ &\leq r \sum_{< \beta, q > = 1} |a_{j\beta}(x) - a_{j\beta}(x_0)| + \sum_{< \beta, q > < 1} r^{< \beta, q >} |a_{j\beta}(x)|, \end{aligned}$$

by the mean-value theorem and the choice of ε , we obtain

$$|P_j(x, r^q \xi_0)| \leq r \sum_{< \beta, q > = 1} C_2 \delta r^{-\varepsilon/\mu} + \sum_{< \beta, q > < 1} C' r^{1-\varepsilon/\mu} \leq C_1 r^{1-\varepsilon/\mu}$$

Since $\varphi \in G_0^{q, \sigma \mu}(\mathbb{R}^n)$, so $\exists C_0 > 0, \exists L_0 > 0, \forall \beta \in \mathbb{Z}_+^n, \forall x \in \mathbb{R}^n$

$$(3.4) \quad |D^\beta \varphi(x)| \leq C_0 L_0^{|\beta|} [\Gamma(< \beta, q > + 1)]^{\sigma \mu}$$

We choose $L_0 \geq 2MC_2^{\sigma \mu}$, where C_2 is the constant of the following inequality $\exists C > 0, \forall \gamma \in \mathbb{Z}_+^n, \forall \beta \in \mathbb{Z}_+^n$,

$$(3.5) \quad \frac{\gamma!}{\beta! (\gamma - \beta)!} \leq C^{< \gamma - \beta, q >} \frac{\Gamma(< \gamma, q > + 1)}{\Gamma(< \beta, q > + 1) \Gamma(< \gamma - \beta, q > + 1)}$$

In the sequel we will use the following properties of the gamma function.

$$(3.6) \quad \Gamma(a + p) = \Gamma(a) a(a + 1) \cdots (a + p - 1), \quad a > 0, p \in \mathbb{Z}_+^*$$

$$\forall \lambda > 0, \forall \tau > 0, \forall a \geq \omega, \forall b \geq \omega, \forall c \geq \omega, \forall \sigma \geq 1$$

$$(3.7) \quad \lambda^a \Gamma(b + c + 1) \sigma \tau^c \leq 2^{\frac{a}{\sigma}} [\lambda^{a+c} \Gamma(b + 1)^\sigma + \Gamma(a + b + c + 1)^\sigma \tau^{a+c}].$$

We need a convenient form of $P_{i_k} \cdots P_{i_1} u$ for any integer $k \geq 0$. By the generalized Leibniz formula $P_j(x, D)(uv) = \sum_{\alpha} \frac{1}{\alpha!} P_j^{(\alpha)} u D^\alpha v$, we obtain

$$\begin{aligned} P_j u(x) &= \int_1^{+\infty} \sum_{< \alpha, q > \leq 1} \frac{1}{\alpha!} D_x^\alpha \left(\varphi[r^{\varepsilon q}(x - x_0)] e^{-r^\eta} \right) P_j^{(\alpha)} \left(e^{i < x - x_0, r^q \xi_0 >} \right) dr \\ &= \int_1^{+\infty} \sum_{< \alpha, q > \leq 1} \frac{1}{\alpha!} P_j^{(\alpha)}(x, r^q \xi_0) D_x^\alpha \left(\varphi[r^{\varepsilon q}(x - x_0)] \right) e^{-r^\eta} e^{i < x - x_0, r^q \xi_0 >} dr \\ &= \int_1^{+\infty} A_j(x, r) e^{-r^\eta} e^{i < x - x_0, r^q \xi_0 >} dr, \end{aligned}$$

where

$$A_j(x, r) = \sum_{< \alpha, q > \leq 1} \frac{1}{\alpha!} P_j^{(\alpha)}(x, r^q \xi_0) D_x^\alpha \left(\varphi[r^{\varepsilon q}(x - x_0)] \right)$$

For any integer $k \geq 0$, $l \leq k$ and $1 \leq i_l \leq N$, we write

$$P_{i_k} \dots P_{i_0} u(x) = \int_1^{+\infty} A_{i_k \dots i_0}(x, r) e^{-r^\eta} e^{i \langle x - x_0, r^q \xi_0 \rangle} dr, \quad ,$$

where P_{i_0} designs the identity operator, and

$$(3.8) \quad \begin{cases} A_{i_0}(x, r) = \varphi[r^{\varepsilon q}(x - x_0)] \\ A_{i_{k+1}, i_k \dots i_0}(x, r) = \sum_{\langle \alpha, q \rangle \leq 1} \frac{1}{\alpha!} P_{i_{k+1}}^{(\alpha)}(x, r^q \xi_0) D_x^\alpha A_{i_k \dots i_0}(x, r) \end{cases}$$

To complete the proof we need the following

LEMMA 4. $\exists L > 0$, $\forall k \in \mathbb{Z}_+$, $\forall \gamma \in \mathbb{Z}_+^n$, $\forall x \in B_0$, $\forall r \geq 1$,

$$(3.9) \quad \begin{aligned} |D_x^\gamma A_{i_k \dots i_0}(x, r)| &\leq C_0 (L_0 r^\varepsilon)^{|\gamma|} L^k \left(r^{(1-\varepsilon/\mu)k} [\Gamma(\langle \gamma, q \rangle + 1)]^{\sigma\mu} \right. \\ &\quad \left. + [\Gamma(\langle \gamma, q \rangle + k + 1)]^{\sigma\mu} r^{k\varepsilon(2-1/\mu)} \right) \end{aligned}$$

Proof. It's obtained by recurrence over k . In fact for $k = 0$, this is true since $\varphi \in G_0^{q, \sigma\mu}(\mathbb{R}^n)$, i.e. we find (3.4). So suppose that the estimate (3.9) holds up to the order k and check that it remains valid at the order $k + 1$. Set

$$\lambda = r^{1-\varepsilon/\mu},$$

$$\tau = r^{\varepsilon(2-1/\mu)},$$

$$S(k, \beta) = \lambda^k [\Gamma(\langle \beta, q \rangle + 1)]^{\sigma\mu} + [\Gamma(\langle \beta, q \rangle + k + 1)]^{\sigma\mu} \tau^k.$$

Then the estimate (3.9) is written as

$$|D_x^\gamma A_{i_k \dots i_0}(x, r)| \leq C_0 (L_0 r^\varepsilon)^{|\gamma|} L^k S(k, \gamma)$$

Let $\omega = \min_{1 \leq j \leq n} q_j$, from inequality (3.7), we have

$$(3.10) \quad r^{1-\varepsilon/\mu} S(k, \beta) \leq 2^{\frac{\sigma\mu}{\omega}+1} S(k+1, \beta),$$

indeed

$$\begin{aligned} r^{1-\varepsilon/\mu} S(k, \beta) &= \lambda \left(\lambda^k [\Gamma(\langle \beta, q \rangle + 1)]^{\sigma\mu} + [\Gamma(\langle \beta, q \rangle + k + 1)]^{\sigma\mu} \tau^k \right) \\ &= \lambda^{k+1} [\Gamma(\langle \beta, q \rangle + 1)]^{\sigma\mu} + \lambda [\Gamma(\langle \beta, q \rangle + k + 1)]^{\sigma\mu} \tau^k \\ &\leq 2^{\frac{\sigma\mu}{\omega}} \left(\lambda^{k+1} [\Gamma(\langle \beta, q \rangle + 1)]^{\sigma\mu} + [\Gamma(\langle \beta, q \rangle + (k+1) + 1)]^{\sigma\mu} \tau^{k+1} \right. \\ &\quad \left. + \lambda^{k+1} [\Gamma(\langle \beta, q \rangle + 1)]^{\sigma\mu} + [\Gamma(\langle \beta, q \rangle + (k+1) + 1)]^{\sigma\mu} \tau^{k+1} \right) \\ &\leq 2^{\frac{\sigma\mu}{\omega}+1} S(k+1, \beta) \end{aligned}$$

Similarly we find

$$(3.11) \quad \lambda^{1-\langle \alpha, q \rangle} \tau^{\langle \alpha, q \rangle} S(k, \beta + \alpha) \leq 2^{\frac{\sigma\mu}{\omega}+1} S(k+1, \beta), \quad \langle \alpha, q \rangle \leq 1$$

From (3.8), we have

$$\begin{aligned} |D_x^\gamma A_{i_{k+1} \dots i_0}(x, r)| &= \left| \sum_{\langle \alpha, q \rangle \leq 1} \frac{1}{\alpha!} D_x^\gamma \left(P_{i_{k+1}}^{(\alpha)}(x, r^q \xi_0) D_x^\alpha A_{i_k \dots i_0}(x, r) \right) \right| \\ &\leq \left| \sum_{\langle \alpha, q \rangle \leq 1} \sum_{\beta \leq \gamma} \frac{1}{\alpha!} \binom{\gamma}{\beta} D_x^{\gamma-\beta} P_{i_{k+1}}^{(\alpha)}(x, r^q \xi_0) D_x^{\alpha+\beta} A_{i_k \dots i_0}(x, r) \right| \\ &\leq I_1 + I_2 + I_3, \end{aligned}$$

where

$$\begin{aligned}
I_1 &= |P_{i_{k+1}}(x, r^q \xi_0)| |D_x^\gamma A_{i_k \dots i_0}(x, r)| \\
I_2 &= \sum_{\beta < \gamma} \binom{\gamma}{\beta} |D_x^{\gamma-\beta} P_{i_{k+1}}(x, r^q \xi_0)| |D_x^\beta A_{i_k \dots i_0}(x, r)| \\
I_3 &= \sum_{0 < \alpha, q \leq 1} \sum_{\beta \leq \gamma} \frac{1}{\alpha!} \binom{\gamma}{\beta} |D_x^{\gamma-\beta} P_{i_{k+1}}^{(\alpha)}(x, r^q \xi_0)| |D_x^{\alpha+\beta} A_{i_k \dots i_0}(x, r)|
\end{aligned}$$

Since the $A_{i_k \dots i_0}$ are functions of compact supports in $B(x_0, 2\delta r^{-\varepsilon/\mu})$, so due to (3.3) and (3.10), we have

$$\begin{aligned}
I_1 &\leq C_1 r^{1-\varepsilon/\mu} C_0 (L_0 r^\varepsilon)^{|\gamma|} S(k, \gamma) L^k \\
(3.12) \quad &\leq 2^{\frac{\sigma\mu}{\omega}+1} C_1 C_0 (L_0 r^\varepsilon)^{|\gamma|} S(k+1, \gamma) L^k
\end{aligned}$$

Using (3.2) and (3.10), we obtain

$$\begin{aligned}
I_2 &\leq \sum_{\beta < \gamma} \binom{\gamma}{\beta} M^{|\gamma-\beta|+1} [\Gamma(<\gamma-\beta, q>+1)]^{\sigma\mu} r C_0 (L_0 r^\varepsilon)^{|\beta|} S(k, \beta) L^k \\
&\leq \sum_{\beta < \gamma} \binom{\gamma}{\beta} [\Gamma(<\gamma-\beta, q>+1)]^{\sigma\mu} M^{|\gamma-\beta|+1} r^\varepsilon C_0 (L_0 r^\varepsilon)^{|\beta|} 2^{\frac{\sigma\mu}{\omega}+1} S(k+1, \beta) L^k
\end{aligned}$$

In the other hand, (3.5) and (3.6), give

$$(3.13) \quad \binom{\gamma}{\beta} [\Gamma(<\gamma-\beta, q>+1)]^{\sigma\mu} S(k, \beta) \leq C_2^{\sigma\mu <\gamma-\beta, q>} S(k, \gamma)$$

Thus we obtain

$$\begin{aligned}
I_2 &\leq \sum_{\beta < \gamma} C_2^{\sigma\mu |\gamma-\beta|} M^{|\gamma-\beta|+1} r^\varepsilon C_0 (L_0 r^\varepsilon)^{|\beta|} 2^{\frac{\sigma\mu}{\omega}+1} S(k+1, \gamma) L^k \\
&\leq \sum_{\beta < \gamma} \left(\frac{M C_2^{\sigma\mu}}{L_0 r^\varepsilon} \right)^{|\gamma-\beta|} r^\varepsilon 2^{\frac{\sigma\mu}{\omega}+1} M C_0 (L_0 r^\varepsilon)^{|\gamma|} S(k+1, \gamma) L^k \\
&\leq \sum_{0 < \beta \leq \gamma} \left(\frac{M C_2^{\sigma\mu}}{L_0 r^\varepsilon} \right)^{|\beta|} r^\varepsilon 2^{\frac{\sigma\mu}{\omega}+1} M C_0 (L_0 r^\varepsilon)^{|\gamma|} S(k+1, \gamma) L^k \\
&\leq \frac{n M C_2^{\sigma\mu}}{L_0 r^\varepsilon} \sum_{\beta \geq 0} \left(\frac{M C_2^{\sigma\mu}}{L_0 r^\varepsilon} \right)^{|\beta|} r^\varepsilon 2^{\frac{\sigma\mu}{\omega}+1} M C_0 (L_0 r^\varepsilon)^{|\gamma|} S(k+1, \gamma) L^k
\end{aligned}$$

Set $C_3 = \sum_{\alpha \geq 0} \left(\frac{1}{2}\right)^{|\alpha|}$, since $L_0 \geq 2M C_2^{\sigma\mu}$ and $r \geq 1$, then

$$(3.14) \quad I_2 \leq \frac{n M C_2^{\sigma\mu}}{L_0} C_3 2^{\frac{\sigma\mu}{\omega}+1} M C_0 (L_0 r^\varepsilon)^{|\gamma|} S(k+1, \gamma) L^k$$

Finally in view of (3.2)

$$\begin{aligned}
I_3 &\leq \sum_{0 < \alpha, q \leq 1} \sum_{\beta \leq \gamma} \binom{\gamma}{\beta} [\Gamma(<\gamma-\beta, q>+1)]^{\sigma\mu} M^{|\gamma-\beta|+1} r^{1-<\alpha, q>} \\
&\quad \times C_0 (L_0 r^\varepsilon)^{|\beta+\alpha|} S(k, \beta+\alpha) L^k
\end{aligned}$$

For any $\alpha \in Z_+^n$, $0 < \alpha, q \leq 1$, we have

$$r^{1-<\alpha, q>+\varepsilon <\alpha, q>} \leq \lambda^{1-<\alpha, q>} \tau^{<\alpha, q>} \quad ,$$

which gives, with (3.11) and (3.13),

$$\begin{aligned} I_3 &\leq \sum_{0 < \alpha, q \leq 1} \sum_{\beta \leq \gamma} \binom{\gamma}{\beta} [\Gamma(\gamma - \beta, q > +1)]^{\sigma\mu} M^{|\gamma - \beta| + 1} \\ &\quad \times C_0 (L_0 r^\varepsilon)^{|\beta|} L_0^{|\alpha|} 2^{\frac{\sigma\mu}{\omega} + 1} S(k+1, \beta) L^k \\ &\leq \sum_{0 < \alpha, q \leq 1} \sum_{\beta \leq \gamma} \left(\frac{MC_2^{\sigma\mu}}{L_0 r^\varepsilon} \right)^{|\gamma|} 2^{\frac{\sigma\mu}{\omega} + 1} MC_0 L_0^{|\alpha|} (L_0 r^\varepsilon)^{|\gamma|} S(k+1, \gamma) L^k \end{aligned}$$

Put $C_4 = \sum_{0 < \alpha, q \leq 1} L_0^{|\alpha|}$, then we obtain

$$(3.15) \quad I_3 \leq 2^{\frac{\sigma\mu}{\omega} + 1} MC_4 C_3 C_0 (L_0 r^\varepsilon)^{|\gamma|} S(k+1, \gamma) L^k$$

If we choose

$$L \geq 2^{\frac{\sigma\mu}{\omega} + 1} \left(C_1 + \frac{nM^2 C_2^{\sigma\mu}}{L_0} C_3 + MC_3 C_4 \right),$$

we get, from (3.12), (3.14) and (3.15),

$$I_1 + I_2 + I_3 \leq C_0 (L_0 r^\varepsilon)^{|\gamma|} S(k+1, \gamma) L^{k+1},$$

which means that (3.9) holds at the order $k+1$.

End of the proof of theorem 3. Applying the last lemma for $\gamma = 0$, we find

$$\begin{aligned} |A_{i_k \dots i_0}(x, r)| &\leq C_0 L^k \left(r^{(1-\varepsilon/\mu)k} + [\Gamma(k+1)]^{\sigma\mu} r^{k\varepsilon(2-1/\mu)} \right) \\ (3.16) \quad &\leq C'_0 L^k \left(r^{(1-\varepsilon/\mu)k} + (k!)^{\sigma\mu} r^{k\varepsilon(2-1/\mu)} \right) \end{aligned}$$

In the other hand, we know that

$$\forall \lambda \in \mathbb{R}_+, \forall k \in \mathbb{Z}_+, \forall s > 0, \quad \frac{\left(\frac{\lambda^{1/s\mu}}{2s\mu} \right)^k}{k!} \leq \exp \left(\frac{\lambda^{1/s\mu}}{2s\mu} \right)$$

So

$$\begin{cases} \lambda^k \leq (2s\mu)^{k\mu s} (k!)^{s\mu} \exp \left(\frac{\lambda^{1/s\mu}}{2} \right) \\ \tau^k \leq (2(s-\sigma)\mu)^{k\mu(s-\sigma)} (k!)^{(s-\sigma)\mu} \exp \left(\frac{\mu^{1/(s-\sigma)\mu}}{2} \right) \end{cases}$$

Thus we obtain, with (3.16),

$$|A_{i_k \dots i_0}(x, r)| \leq C'_0 L^k (2s\mu)^{k\mu s} (k!)^{s\mu} \left[\exp \left(\frac{r^\eta}{2} \right) + \exp \left(\frac{r^{\eta'}}{2} \right) \right],$$

where

$$\eta' = \frac{\varepsilon(2-1/\mu)}{\mu(s-\sigma)} \leq \eta = \frac{1-\varepsilon/\mu}{\mu s},$$

since $\varepsilon \leq \frac{\mu(s-\sigma)}{2\mu s - \sigma}$. Therefore

$$|A_{i_k \dots i_0}(x, r)| \leq 2C'_0 L^k (k!)^{s\mu} \exp \left(\frac{r^\eta}{2} \right)$$

The last estimate gives

$$\begin{aligned} |P_{i_k \dots i_0} u(x)| &\leq 2C'_0 L'^k (k!)^{s\mu} \int_1^{+\infty} \exp\left(-\frac{r^\eta}{2}\right) dr \\ &\leq C^{k+1} (k!)^{s\mu}, \end{aligned}$$

which means that $u \in G^s\left(\Omega, (P_j)_{j=1}^N\right)$ ■

4. Consequences

The theorems of this work give and unify the results of Bolley-Camus [1] and Métivier [6] in the homogenous case, and the results of Bouzar-Chaïli [3] in the quasi-homogenous case.

COROLLARY 2. *Let Ω be an open subset of $\mathbb{R}^n$ and $\sigma > s \geq 1$, and let $(P_j)_{j=1}^N$ be a system of linear differential operators with coefficients in $G^{q,\sigma}(\Omega)$, then*

$$(P_j)_{j=1}^N \text{ is } q\text{-quasi-elliptic in } \Omega \iff G^s\left(\Omega, (P_j)_{j=1}^N\right) \subset G^{q,s}(\Omega)$$

Theorem 2 is also a generalization of the main result of [7].

COROLLARY 3. *Let Ω be an open subset of $\mathbb{R}^n$, $s \geq 1$ and let P be a linear differential operator with coefficients in $G^{0,s}(\Omega)$, then*

$$P \text{ is multi-quasi-elliptic in } \Omega \Rightarrow G^s(\Omega, P) \subset G^{\mathcal{F},s}(\Omega)$$

REMARK 2. *In [7] the necessity has not been given even for the scalar case.*

References

- [1] P. Bolley, J. Camus, Powers and Gevrey regularity for a system of differential operators, Czechoslovak Math. J., Praha, 29 (104), (1979), 649-661.
- [2] P. Bolley, J. Camus, L. Rodino, Hypoellipticité analytique-Gevrey et itérés d'opérateurs, Ren. Sem. Mat. Univ. Politec. Torino, vol. 45:3, (1988), 1-61.
- [3] C. Bouzar, R. Chaïli, Vecteurs Gevrey d'opérateurs différentiels quasihomogènes, to appear in Bul. Belg Math. Soc. Simon Stevin.
- [4] C. Bouzar, R. Chaïli, Une généralisation de la propriété des itérés, Arch. Math., 76:1, (2001), 57-66.
- [5] S.G. Gindikin, L.R. Volevich, The method of Newton's polyhedron in the theory of partial differential equations, Kluwer Acad. Publ., 1992.
- [6] G. Métivier, Propriété des itérés et ellipticité, Comm. Partial. Differential Equations, 3, 9, (1978), 827-876.
- [7] L. Zanghirati, Iterati di una classe di operatori ipoellittici e classi generalizzate di Gevrey, Boll. U.M.I., vol. 1, suppl., (1980), 177-195.

DÉPARTEMENT DE MATHÉMATIQUES. UNIVERSITÉ D'ORAN ESENIA. ALGÉRIE.
E-mail address: bouzarchikh@hotmail.com